arXiv:1607.07814v2 [math.CO] 10 Mar 2017

MINKOWSKI COMPLEXES AND CONVEX THRESHOLD DIMENSION

FLORIAN FRICK AND RAMAN SANYAL

Abstract. For a collection of convex bodies $P_1, \dots, P_n \subset \mathbb{R}^d$ containing the origin, a *Minkowski complex* is given by those subsets whose Minkowski sum does not contain a fixed basepoint. Every simplicial complex can be realized as a Minkowski complex and for convex bodies on the real line, this recovers the class of threshold complexes. The purpose of this note is the study of the *convex threshold dimension* of a complex, that is, the smallest dimension in which it can be realized as a Minkowski complex. In particular, we show that the convex threshold dimension can be arbitrarily large. This is related to work of Chvátal and Hammer (1977) regarding forbidden subgraphs of threshold graphs. We also show that convexity is crucial this context.

A simplicial complex Δ on vertices $[n] := \{1, \dots, n\}$ is a **threshold complex** if there are real numbers $\lambda_1, \dots, \lambda_n, \mu \in \mathbb{R}$ with $0 \le \lambda_i \le \mu$ for all $i = 1, \dots, n$ such that for any $\sigma \subseteq [n]$

$$\sigma \ \in \ \Delta \qquad \text{if and only if} \qquad \sum_{i \in \sigma} \lambda_i \ < \ \mu.$$

Threshold complexes (or hypergraphs) where proposed by Golumbic [4] as a higher-dimensional generalization of the threshold *graphs* of Chvátal and Hammer [2]; see also [9]. If we assume that $0 \le \lambda_1 \le \cdots \le \lambda_n \le \mu$, then for any $i \in \sigma \in \Delta$ and $j < i$, we have $(\sigma \setminus i) \cup j \in \Delta$. Hence, threshold complexes are **shifted** in the sense of Kalai [6] and topologically wedges of (not necessarily equidimensional) spheres. See [7] and [3] for more information regarding the combinatorics and topology of threshold and shifted complexes.

The purpose of this note is to investigate a generalization of threshold complexes inspired by convex geometry. For that, let $\mathcal{P} = (P_1, \dots, P_n)$ be an ordered family of convex bodies in $\mathbb{R}^d$ each containing the origin and let $\mu \in \mathbb{R}^d$ be a point. The **Minkowski complex** associated to $\mathcal{P}$ and μ is the simplicial complex $\Delta(\mathcal{P}; \mu)$ given by the simplices $\sigma \subseteq [n]$ with

$$\sigma \ \in \ \Delta(\mathcal{P}; \mu) \qquad \text{if and only if} \qquad \mu \ \notin \ P_\sigma \ := \ \sum_{i \in \sigma} P_i.$$

Here, $\sum_{i \in \sigma} P_i \ = \ \{\sum_{i \in \sigma} p_i : p_i \in P_i\}$ is the **Minkowski sum** (or vector sum) and we set $P_\emptyset := \{0\}$. By setting $P_i := \{t \in \mathbb{R} : 0 \le t \le \lambda_i\}$, it follows that threshold complexes are Minkowski complexes. For the case that each $P_i \subset \mathbb{R}^d$ is an axis-parallel box, these simplicial complexes have been studied by Pakianathan and Winfree [8] under the name of *quota complexes*. We may also replace a convex body P_i by a suitable convex polytope and we will tacitly do this henceforth.

Our motivation for studying Minkowski complexes comes from *mixed Ehrhart theory*. For a set $S \subset \mathbb{R}^d$, let us define the **discrete volume** $E(S) := |S \cap \mathbb{Z}^d|$. The **discrete mixed volume**

Date: March 13, 2017.

2010 *Mathematics Subject Classification.* 05E45, 52A35, 52A39.

Key words and phrases. Minkowski sum, Minkowski complexes, threshold complexes, convex threshold dimension, discrete mixed volume.

of lattice polytopes $P_1, \dots, P_n \subset \mathbb{R}^d$ is defined as

$$\mathrm{CM}E(P_1, \dots, P_n) \;=\; \sum_{I \subseteq [n]} (-1)^{n-|I|} E(P_I).$$

It was shown in [5] (see also [1]) that, like its continuous counterpart the *mixed volume*, the discrete mixed volume satisfies

$$0 \;\le\; \mathrm{CM}E(Q_1, \dots, Q_n) \;\le\; \mathrm{CM}E(P_1, \dots, P_n)$$

for lattice polytopes $Q_i \subseteq P_i$ for $i = 1, \dots, n$. Since $\mathrm{CM}E$ is invariant under lattice translations, we may assume that $0 \in P_i$ for all i. This allows us to express the discrete mixed volume as follows:

Theorem 1. *Let $\mathcal{P} = (P_1, \dots, P_n)$ be a family of $n > 0$ lattice polytopes in $\mathbb{R}^d$ with $0 \in P_i$ for all i. Then*

$$(-1)^n \mathrm{CM}E(P_1, \dots, P_n) \;=\; \sum_{\mu \in P_{[n]} \cap \mathbb{Z}^d} \tilde{\chi}(\Delta(\mathcal{P}; \mu)),$$

where $\tilde{\chi}$ denotes the reduced Euler characteristic.

Proof. Since all polytopes P_i contain the origin, it follows that $P_I \subseteq P_J$ for $I \subseteq J$. Let us write $[P_I] : \mathbb{R}^d \to \{0, 1\}$ for the characteristic function of P_I and define $F := \sum_I (-1)^{n-|I|} [P_I]$. Then $\mathrm{CM}E(P_1, \dots, P_n) = \sum_{\mu \in \mathbb{Z}^d} F(\mu)$. The result now follows from the observation that

$$(-1)^n F(\mu) \;=\; \sum_{\sigma \subseteq [n],\, \mu \notin P_\sigma} (-1)^{|\sigma|} \;=\; \tilde{\chi}(\Delta(\mathcal{P}, \mu))$$

for all $\mu \in \mathbb{R}^d$. □

Theorem 1 prompted the question if Minkowski complexes have restricted topology. It was already noted in [8, Thm. D.1] that this is not the case. We recall their result with a short proof.

Proposition 2. *For any simplicial complex $\Delta \subseteq 2^{[n]}$ there are polytopes $\mathcal{P} = (P_1, \dots, P_n)$ and μ in some $\mathbb{R}^D$ such that $\Delta = \Delta(\mathcal{P}; \mu)$.*

Proof. Let $\sigma_1, \dots, \sigma_D$ be the facets of Δ. For $1 \le i \le n$ and $1 \le j \le D$ we set $t_{ij} := 1$ if $i \in \sigma_j$ and $t_{ij} := |\sigma_j| + 1$ otherwise. For $1 \le i \le n$ define

$$P_i \;:=\; \{p \in \mathbb{R}^D : 0 \le p_j \le t_{ij} \text{ for } 1 \le j \le D\}$$

and let $\mu = (|\sigma_1| + 1, \dots, |\sigma_d| + 1)$. Observe that $\mu \notin \sum_{i \in \sigma_j} P_i$ for all j and thus $\Delta \subseteq \Delta(\mathcal{P}; \mu)$. Conversely, if $\tau \notin \Delta$, then for any j there is an i with $i \in \tau \setminus \sigma_j$. This implies $\mu \in P_\tau$. □

As a measure of complexity, define the **convex threshold dimension** $\mathsf{ctd}(\Delta)$ of a simplicial complex $\Delta \subseteq 2^{[n]}$ as smallest dimension d in which Δ can be realized as a Minkowski complex. This is exactly the minimum over $\dim P_1 + \cdots + P_n$ over all $(\mathcal{P}, \mu)$ for which $\Delta = \Delta(\mathcal{P}; \mu)$. Thus, the empty complex is the unique complex of convex threshold dimension 0 while $\mathsf{ctd}(\Delta) = 1$ if and only if Δ is a threshold complex. Proposition 2 shows that the convex threshold dimension is finite for every simplicial complex. In the remainder we will show that $\mathsf{ctd}(\Delta)$ can be arbitrarily large and hence is a *proper* measure of the complexity of Δ.

In [2] it was shown that threshold graphs are characterized by the three forbidden induced subgraphs given in Figure 1. The following result shows that these graphs are key to increasing



FIGURE 1. The three forbidden induced subgraphs for threshold graphs.

the convex threshold dimension. Let us denote by $\Delta * \Gamma = \{\sigma \uplus \tau : \sigma \in \Delta, \tau \in \Gamma\}$ the join of simplicial complexes Δ and Γ.

Theorem 3. *Let Γ be a simplicial complex that contains one of the graphs of Figure 1 as an induced subgraph. Then*

$$\mathsf{ctd}(\Delta * \Gamma) \;\geq\; \mathsf{ctd}(\Delta) + 1$$

for every simplicial complex Δ.

Corollary 4. *For every $d \geq 0$, there is a simplicial complex Δ with $\mathsf{ctd}(\Delta) \geq d$.*

For the proof of Theorem 3, we first note the following two helpful facts about Minkowski complexes.

Lemma 5. *Let $\Delta = \Delta(\mathcal{P}; \mu)$ be a Minkowski complex for $P_1, \ldots, P_n \subset \mathbb{R}^d$. Let $\sigma, \tau \in \Delta$ be faces such that $\sigma \cup \tau \in \Delta$. For any line ℓ through μ, then the restrictions of P_σ and P_τ to ℓ are on the same side of μ.*

Proof. Since $\sigma \cup \tau \in \Delta$ the convex set $P_{\sigma\cup\tau} = P_\sigma + P_\tau$ does not contain μ. By assumption on $\mathcal{P}$, we have $P_\sigma \cup P_\tau \subseteq P_{\sigma\cup\tau}$ from which the claim follows. □

Lemma 6. *Let $\Delta = \Delta(\mathcal{P}; \mu)$ be a Minkowski complex for a collection of polytopes $\mathcal{P}$ in $\mathbb{R}^d$. Suppose that there is a line ℓ through μ that does not intersect any P_σ for $\sigma \in \Delta$, then $\mathsf{ctd}(\Delta) < d$.*

Proof. Denote by $\pi\colon \mathbb{R}^d \to \ell^\perp \cong \mathbb{R}^{d-1}$ the orthogonal projection along the line ℓ. Let $\pi(\mathcal{P}) = (\pi(P_1), \ldots, \pi(P_n))$. For $\sigma \in \Delta$ the set P_σ avoids ℓ and thus $\pi(\mu) \notin \pi(P_\sigma)$. This shows that $\sigma \in \Delta(\pi(\mathcal{P}); \pi(\mu))$. Conversely $\sigma \in \Delta(\pi(\mathcal{P}); \pi(\mu))$ implies that the line $\ell = \pi^{-1}(\mu)$ does not intersect P_σ and thus $\sigma \in \Delta(\mathcal{P}; \mu)$. The claim now follows from $\dim \pi(P_1) + \cdots + \pi(P_n) < d$. □

Proof of Theorem 3. Suppose that $\mathsf{ctd}(\Delta * \Gamma) = d$ and fix a realization $\Delta(\mathcal{P}; \mu)$ with $\mathcal{P}$ a collection of polytopes in $\mathbb{R}^d$. Denote the two edges of Γ that induce one of the graphs in Figure 1 by e and f and let e' and f' be two disjoint (diagonal) nonedges. Then $\mu \in P_{e'} \cap P_{f'}$. Since $e \cup f = e' \cup f'$ we have $2\mu \in P_{e'} + P_{f'} = P_e + P_f$. Thus, there is a vector $v \in \mathbb{R}^d \setminus \{0\}$ with $\mu - v \in P_e$ and $\mu + v \in P_f$. The line ℓ connecting $\mu - v$ and $\mu + v$ goes through μ. The convex sets P_e and P_f intersect ℓ on different sides of μ.

The line ℓ must also intersect P_σ for some face $\sigma \in \Delta$ by Lemma 6. Since $\sigma \cup e, \sigma \cup f \in \Delta * \Gamma$, the sets P_σ and P_e as well as P_σ and P_f intersect ℓ on the same side of μ by Lemma 5. This is a contradiction. □

It would be very interesting if complexes of convex threshold dimension d can be characterized in terms of the number of distinct copies of the forbidden subgraphs.

As a last thought, let us emphasize that convexity played a crucial role in our considerations. For that, observe that the definition of Minkowski complex $\Delta(\mathcal{X}; \mu)$ makes sense for collections $\mathcal{X} = (X_1, \ldots, X_n)$ of *arbitrary* sets in $\mathbb{R}^d$ that contain the origin.

Proposition 7. *Let Δ be a simplicial complex. Then there is a collection $\mathcal{X} = (X_1, \dots, X_n)$ of discrete sets in $\mathbb{R}$ such that $\Delta = \Delta(\mathcal{X}; \mu)$ for some $\mu \in \mathbb{R}$. In particular, there is a collection $\mathcal{X}$ of contractible sets in $\mathbb{R}^2$ realizing Δ as a Minkowski complex.*

Proof. Let $(\mathcal{P}; \mu)$ be a realization of Δ by convex sets in some $\mathbb{R}^d$. For any $\tau \notin \Delta$, choose $y_i^\tau \in P_i$ such that $\mu = y_1^\tau + \cdots + y_n^\tau$. Then $Y_i := \{0\} \cup \{y_i^\tau : \tau \notin \Delta\}$ yields a realization of Δ by discrete sets in $\mathbb{R}^d$. The argument in the proof of Lemma 6 applies unless $d = 1$ and proves the first claim. The second claim simply follows from the fact that we may connect the points $x_i \in X_i \setminus \{0\}$ by internally disjoint arcs properly contained in the upper half-plane to 0. □

References

[1] F. Bihan, *Irrational Mixed Decomposition and Sharp Fewnomial Bounds for Tropical Polynomial Systems*, Discrete Comput. Geom., 55 (2016), pp. 907–933. 2

[2] V. Chvátal and P. L. Hammer, *Aggregation of inequalities in integer programming*, in Studies in integer programming (Proc. Workshop, Bonn, 1975), North-Holland, Amsterdam, 1977, pp. 145–162. Ann. of Discrete Math., Vol. 1. 1, 2

[3] P. H. Edelman, T. Gvozdeva, and A. Slinko, *Simplicial complexes obtained from qualitative probability orders*, SIAM J. Discrete Math., 27 (2013), pp. 1820–1843. 1

[4] M. C. Golumbic, *Algorithmic graph theory and perfect graphs*, Academic Press [Harcourt Brace Jovanovich, Publishers], New York-London-Toronto, Ont., 1980. With a foreword by Claude Berge, Computer Science and Applied Mathematics. 1

[5] K. Jochemko and R. Sanyal, *Combinatorial mixed valuations*. Preprint, May 2016, 16 pages, arXiv:1605.07431. 2

[6] G. Kalai, *Algebraic shifting*, in Computational commutative algebra and combinatorics (Osaka, 1999), vol. 33 of Adv. Stud. Pure Math., Math. Soc. Japan, Tokyo, 2002, pp. 121–163. 1

[7] C. J. Klivans, *Threshold graphs, shifted complexes, and graphical complexes*, Discrete Math., 307 (2007), pp. 2591–2597. 1

[8] J. Pakianathan and T. Winfree, *Threshold complexes and connections to number theory*, Turkish J. Math., 37 (2013), pp. 511–539. 1, 2

[9] J. Reiterman, V. Rödl, E. Šiňajová, and M. Tuma, *Threshold hypergraphs*, Discrete Math., 54 (1985), pp. 193–200. 1

Department of Mathematics, Cornell University, Ithaca, NY 14853, USA

E-mail address: ff238@cornell.edu

Institut für Mathematik, Goethe-Universität Frankfurt, Germany

E-mail address: sanyal@math.uni-frankfurt.de